

Physical Constraints and Functional Characteristics of Transcription Factor-DNA Interaction

Ulrich Gerland*, J. David Moroz[†]*, and Terence Hwa

Department of Physics, University of California at San Diego, La Jolla, CA 92093-0319

(Dated: February 2, 2008)

We study theoretical “design principles” for transcription factor-DNA interaction in bacteria, focusing particularly on the statistical interaction of the transcription factors (TF’s) with the genomic background (i.e., the genome without the target sites). We introduce and motivate the concept of *programmability*, i.e. the ability to set the threshold concentration for TF binding over a wide range merely by mutating the binding sequence of a target site. This functional demand, together with physical constraints arising from the thermodynamics and kinetics of TF-DNA interaction, leads us to a narrow range of “optimal” interaction parameters. We find that this parameter set agrees well with experimental data for the interaction parameters of a few exemplary prokaryotic TF’s. This indicates that TF-DNA interaction is indeed programmable. We suggest further experiments to test whether this is a general feature for a large class of TF’s.

With rapid advances in the sequencing and annotation of entire genomes, the task of understanding the associated regulatory networks becomes increasingly prominent. Currently, many experimental and computational efforts are devoted to deciphering the genetic wiring diagram of a cell [1, 2, 3]. Most of these efforts are focused on locating the functional DNA binding sites of transcription factors (TF’s). This knowledge, together with the genomic sequences, will provide a qualitative picture of which gene products may directly affect the expression of which genes. While obtaining such wiring diagrams is tremendously important for the eventual understanding of gene regulation at the system level, this knowledge in itself is not sufficient for the quantitative understanding of system-level effects. This has been dramatically shown in a detailed experimental study of the regulation of the *endo16* gene in Sea Urchin development [4], which revealed an intricate regulatory function where a dozen or so TF’s control the expression of a *single* gene. It would have been impossible to infer even the gross qualitative features of the transcriptional control from the knowledge of the binding sites alone.

A major obstacle to progress is the lack of a *quantitative* understanding of the physical interaction between the TF’s. However, even the simpler interaction between TF’s and DNA sequences is not so well-understood quantitatively: It is common to classify a potential TF-binding DNA sequence in a “digital” manner — either the sequence is designated for TF-binding or it is not. In this view of TF-DNA interaction, differences between the TF-binding sequences are only nuisances which impede straightforward bioinformatic methods of target sequence discovery. On the other hand, there are plenty of

examples where differences between target sequences are known to be *functionally important* [5]. In many cases, the binding of a TF to one site occurs only in the presence of some other TF, while the binding of the same TF to a different site does not require other TF’s. This is often accomplished by differences in the binding sequences, and is believed to be the basis for combinatorial control and signal integration in gene regulation [6]. Also, different binding sites of the same TF can be “tuned” to bind at different TF concentrations, as suggested by a recent study of the *E. coli* flagella assembly system [7]. If further experimental studies confirm that tuning of binding thresholds is indeed used genome wide to establish desired gene regulatory functions, then TF-DNA binding should be regarded more in an “analog” instead of a “digital” manner.

In this work, we report our theoretical study on the “design” of TF-DNA interaction, assuming the analog scheme of operation. Specifically, we impose the functional requirement that the threshold concentration for TF binding to a site can be controlled over a wide range by the choice of the sequence alone; we refer to this as the “programmability” of TF-DNA binding. Taken together with thermodynamic and kinetic constraints, this functional requirement leads to a narrow range of “optimal” TF-DNA interaction parameters. We then compare our result to experimentally known parameters for exemplary TF’s to determine whether the design of these TF’s would indeed *allow* the analog scheme of operation.

To focus our discussion, we limit ourselves exclusively to the case of bacterial TF’s which are the best characterized experimentally. We study both the equilibrium occupancy of a target sequence, and the dynamics of locating the target. Von Hippel, Berg, and Winter have already discussed many aspects of these issues in a series of seminal articles [8, 9, 10, 11, 12]. Our study is built firmly upon their work, but includes a number of additional issues: (i) the effect of sequence-specific binding to the genomic background (non-target sequences)

*These authors contributed equally to this work.

[†]Present address: NEC Research Institute, 4 Independence Way, Princeton, NJ 08540

on the equilibrium occupation of a target sequence; (ii) kinetic traps arising statistically from the genomic background, and (iii) the desired programmability of TF-DNA binding. We adopt the model developed by von Hippel and Berg, and allow both the sequence-specific and non-specific modes of TF-DNA binding. Sequence-specific binding occurs if the binding sequence is sufficiently close to the best binding sequence, and is governed quantitatively by a specificity parameter. For typical bacterial TF's whose binding sequences are no more than 15 bases long, we find that our physical and functional requirements are best satisfied within a narrow regime of *intermediate specificity*, amounting to the loss of about $2k_B T$ for each additional base mismatch from the best binding sequence. Furthermore, the kinetic constraint favors a low threshold to non-specific binding, while the programmability requirement pushes the threshold to larger values. The optimal trade-off value depends only on the genome size, and lies about $16k_B T$ above the energy of the best binding sequence for a genome of 10^7 bases. These values correspond well with the interaction parameters of a number of well-characterized TF's, which suggests that programmability of TF-DNA binding is compatible with the reality of protein-DNA interaction and may be used by the organism to accomplish biological functions. We hope to stimulate further experiments determining the interaction parameters for a wider range of TF's (see 'Discussion'). These experiments could either strengthen or falsify the programmability concept, depending on whether the interaction parameters are generally in agreement with our prediction.

Model of Transcription Factor-DNA Interaction

Much of our knowledge on the details of TF-DNA interaction is derived from extensive biochemical experiments on a few exemplary systems, dating back to pioneering work in the late 70's [8, 9, 10, 13, 14, 15] and continuing through recent years [16, 17, 18, 19, 20]. Furthermore, detailed structural information is available for many TF's from various structural families [21]. Based on this knowledge, quantitative models of TF-DNA interaction have been established [8, 11, 12, 17]. Together with the recent availability of genomic sequences, these models can be used to characterize the thermodynamics as well as the dynamics of TF's with genomic DNA in a cell. We briefly review the primary model of TF-DNA interaction in this section, which serves to introduce our notation and formulate the problem.

Biochemical and structural experiments, e.g. using *lac*-repressor [9, 14, 20], have established firmly that (i) TF's bind closely to the DNA with a free energy ΔG_{ns} (with respect to the cytoplasm) regardless of its sequence due to electrostatic interaction alone, and (ii) additional sequence-specific binding energy can be gained (via hydrogen bonds) if the binding sequence is close to the recognition sequence of the TF. Let the total binding

(free) energy of a TF to a sequence $\vec{s} = \{s_1, s_2, \dots, s_L\}$ of L nucleotides $s_i \in \{\text{A, C, G, T}\}$ be $\Delta G[\vec{s}]$ (with respect to the cytoplasm), and let $\vec{s}^*$ be the best binding sequence. $\Delta G[\vec{s}]$ becomes sequence-independent, $\Delta G[\vec{s}] \equiv \Delta G_{\text{ns}}$, if $\vec{s}$ is far from $\vec{s}^*$. This is believed to occur via a change in the conformation of the TF, from one which allows more hydrogen bond formation to another which brings the positive charges of the TF closer to the negatively charged DNA backbone [10].

For this study, it will be convenient to measure all energies with respect to that of the best binder, $\Delta G[\vec{s}^*]$. Let us define $E[\vec{s}] \equiv \Delta G[\vec{s}] - \Delta G[\vec{s}^*]$. Furthermore, we will introduce the threshold energy $E_{\text{ns}} \equiv \Delta G_{\text{ns}} - \Delta G[\vec{s}^*]$ where TF-DNA binding switches from the specific to the non-specific mode (for *lac*-repressor, $E_{\text{ns}} \approx 10 \text{ kcal/mole}$). Then given the above model of TF-DNA interaction, and assuming that the TF is bound to the DNA essentially all the time¹, all thermodynamic quantities regarding this TF can be computed from the partition function²

$$Z = \sum_{j=1}^N e^{-\beta E[\vec{s}_j]} + N \cdot e^{-\beta E_{\text{ns}}}, \quad (1)$$

where $\beta^{-1} = k_B T \approx 0.6 \text{ kcal/mole}$ and $\vec{s}_j$ denotes the subsequence of the genomic sequence $\{s_1, s_2, \dots, s_N\}$ from position j to $j + L - 1$. The binding length of a typical bacterial TF is $L = 10 \sim 20$ basepairs (bp). The length of genomic sequence N is typically several million bp.

The form of the binding energy $E[\vec{s}]$ has been studied experimentally for several TF's [16, 17, 18, 19]. In particular, recent experiments on the TF Mnt from bacteriophage P22 [16] support the earlier model [11] that the contribution of each nucleotide in the binding sequence to the total binding energy is approximately independent and additive, i.e.,

$$E[\vec{s}] = \sum_{i=1}^L \mathcal{E}_i(s_i). \quad (2)$$

For the TF's Mnt, Cro, and λ -repressor, the parameters of the "energy matrix" $\mathcal{E}_i(s_i)$ have actually been determined experimentally by *in vitro* measurements of the equilibrium binding constants $K[\vec{s}] \propto e^{-\beta E[\vec{s}]}$ for every *single-nucleotide* mutant of the best binding sequence $\vec{s}^*$ [16, 18, 19]. Due to our definition of the energy scale,

¹ *In vivo* measurements for the case of *lac*-repressor found less than 10% of the TF's were unbound [15]. This agrees well with an estimate based on a typical prokaryotic cell volume of $3 \mu\text{m}^3$, a genome length of $5 \cdot 10^6$ bases, and a non-specific binding constant on the order of 10^4 M^{-1} under physiological conditions [13], which yields a fraction of unbound TF's at a few percent level.

² One should also include the reverse-complement of the genomic sequence in the evaluation of the partition function Z . In order not to make the notation too complicated, we extend the definition of "genomic sequence" to include its complement.

$\mathcal{E}_i(s_i) = 0$ for $s_i = s_i^*$ and $\mathcal{E}_i(s_i) > 0$ for $s_i \neq s_i^*$; the latter will be referred to as “mismatch energies”. While the simple form of the binding energy (2) will certainly not hold for all TF’s, and di-, tri-nucleotide correlation effects are likely to be important in many cases (e.g., to some extent for *lac*-repressor [20]), the key results of our study are not sensitive to such correlations as long as there is a wide range of binding energies for different binding sequences. Thus we will adopt the simple form (2) for this study. For the three well-studied TF’s, the mismatch energies are typically in the range of $1 \sim 3 k_B T$ ’s. While the threshold energies E_{ns} have not been carefully measured for these TF’s, it is believed that non-specific binding does not occur until the binding sequences are at least 4 to 5 mismatches away from $\bar{s}^*$ (G. Stormo, private communication).

Genomic Background and Target Recognition

Thermodynamics

Let us first consider the binding of a *single* TF to its target sequence, denoted by $\bar{s}_t$. We will assume that thermal equilibrium can be reached within the relevant cellular time scale and discuss the important kinetics issue afterwards. The effectiveness of the binding of the TF to its target is then described by the equilibrium binding probability P_t , which depends not only on the binding energy $E_t \equiv E[\bar{s}_t]$, but also on the interaction with the rest of the genomic sequence. Let the contribution of this genomic background to the partition function be Z_b , then the binding probability to the target is given by

$$P_t = \frac{1}{1 + e^{\beta(E_t - F_b)}} , \quad (3)$$

where $F_b = -k_B T \ln Z_b$ is the effective binding energy (or free energy) of the entire genomic background. Eq. (3) is a sigmoidal function of E_t with a (soft) threshold at F_b , i.e., a TF binds (with probability $P_t > 0.5$) if $E_t < F_b$. Since $E_t \geq 0$ by definition, we must have

$$F_b \geq 0 \quad (4)$$

in order for a target sequence to be recognized by a single TF (we consider multiple TF’s below).

The background contribution can be computed for any given TF and genome according to Eq. (1) if the binding energy matrix, the threshold energy E_{ns} , and the genomic sequence is known. We will instead seek a description that is independent of the specifics of the genomic sequences and energy matrices. To accomplish this, we observe first that for the few well-studied TF’s, the interaction of the TF with the genomic background can be well approximated by the interaction of the TF with *random* nucleotide sequences of the same length and single-nucleotide frequencies $p(s)$. This is illustrated in Fig. 1(a) where the histogram of binding energies obtained by using the binding energy matrix $\mathcal{E}_i(s)$ for the




FIG. 1 For the purpose of TF binding, the genome may be treated as random DNA plus functional target site(s): (a) Histogram of the specific binding energies for Cro [solid line] on the *E. coli* genome, together with the average histogram [circles] for Cro on random nucleotide sequences (synthesized with the same length and single-nucleotide frequencies as the *E. coli* genome; normalization for both histograms such that maximum is at N). Except for statistical fluctuations at the low energy end, the histograms are indistinguishable from each other. The approximate position of the threshold energy for nonspecific binding E_{ns} is indicated as the thin straight line. (b) Energy landscape for Cro on the bacteriophage λ DNA. The landscape appears to be *random*, e.g. no “funnel” guides the TF to the target site. The spatial correlation function of the landscape (not shown) decays quickly to zero beyond the scale of $L = 17$ for this case. Random energy landscapes are also found for the other two TF’s with known energy matrices (not shown).

TF Cro on the *E. coli* genome (solid line) coincides well with the histogram of the same energy matrix applied to random nucleotide sequences (circles). Moreover, there appears to be hardly any positional correlation in the binding energies along the genome, as shown by the “energy landscape” in Fig. 1(b) (see caption for details). In the following, we will therefore describe the effect of the genomic background by treating it as a random nucleotide sequence for a generic TF. In particular, we will describe the genomic background partition function by $Z_b = Z_{sp} + N \cdot e^{-\beta E_{ns}}$ where the contribution due to sequence-specific binding is

$$Z_{sp} = \sum_{\bar{s} \in \mathcal{S}(N)} e^{-\beta E[\bar{s}]}, \quad (5)$$

with $\mathcal{S}(N)$ denoting a given collection of N random nucleotide sequences of length L , drawn according to the frequency $p(s)$ for each nucleotide s .

Even with the random sequence approximation (5), computation of the background energy $F_b = -k_B T \ln Z_b$ is nontrivial in principle: From its definition, it is clear that F_b is a random variable, and its precise value will depend on the actual collection of sequences $\mathcal{S}(N)$. We are interested in the *typical* value of F_b , a reasonable approximation of which is its statistical average, $\overline{F_b} \equiv -k_B T \ln \overline{Z_b}$. [We use an overbar to denote averages over an ensemble of different sequence collections $\mathcal{S}(N)$.]

Computing the average $\overline{\ln Z_b}$ is however difficult to do for an arbitrary energy matrix $\mathcal{E}_i(s)$ short of performing numerical simulations. An alternative is to compute the ensemble average of Z_b , i.e., $\overline{Z_b} = \overline{Z_{sp}} + N e^{-\beta E_{ns}}$, where

$$\overline{Z_{sp}} = N \prod_{i=1}^L \left[\sum_{s=\{A,C,G,T\}} e^{-\beta \mathcal{E}_i(s)} p(s) \right] \quad (6)$$

with the single-nucleotide frequencies $p(s)$, and assume that

$$F_b \approx -k_B T \ln \overline{Z_b}. \quad (7)$$

This is, for example, the approach taken by Stormo and Fields [17] in their analysis of the TF Mnt³. We note in passing that $\overline{Z_b}$ can be more compactly written in terms of the density of states $\Omega_{sp}(E)$ for specific binding [the normalized version of the histogram in Fig. 1(a)], i.e.,

$$\overline{Z_b} = N \cdot \sum_E \Omega_{sp}(E) e^{-\beta E} + N \cdot e^{-\beta E_{ns}}. \quad (8)$$

The relation (7) is based on the so-called ‘‘annealed approximation’’ $\overline{\ln Z} \approx \ln \overline{Z}$ which is valid for the genomic sequence length $N \rightarrow \infty$ but not always appropriate for finite N , e.g., if the partition function is dominated by a few low energy terms. Much is known from statistical physics about systems of the type defined by the partition function Z_{sp} in (5), generically known as the Random Energy Model or REM⁴, introduced by Derrida [22]. It turns out that the annealed approximation is valid as long as the system’s entropy S is significantly larger than zero, reflecting the contribution of many terms in the partition sum. We will see further below that proper function of the TF’s requires the system to be in a regime where the annealed approximation is safely applicable. We will thus take the relation (7) for granted. In this case, the condition (4) for the recognition of the target sequence by a single TF becomes

$$\overline{Z_b} \leq 1. \quad (9)$$

Search Dynamics

In order to carry out their function properly, TF’s not only need to have a high equilibrium binding probability to their targets, but also must be able to locate them in a reasonably short time (e.g. less than a few minutes)

after they have been activated by an inducer or freshly produced by a ribosome. This constitutes a constraint on the ‘‘search dynamics’’ of TF’s.

In their non-specific binding mode, TF’s are still strongly associated with the DNA, but are able to diffuse (i.e., slide) randomly along the genome [8, 9, 10]. However, pure one-dimensional diffusion would be an inefficient search process, since it is very redundant (e.g., a 1D random walker always returns back to the start.) For instance, assuming generously a 1D diffusion constant of $D_1 \approx 1 \mu\text{m}^2/\text{sec}$ [10], one finds a time $T_{1D} \sim N^2/D_1 \sim 10^6 \text{ sec}$ for a single TF to diffuse around a bacterial genome of length $N \approx 5 \times 10^6 \text{ bp}$ (about 1 mm). Thus, in order to find a target within a few minutes via 1D diffusion, one would need at least 100 TF’s per cell to search in parallel (so that the search length N is reduced by a factor of 100). On the other hand, there are well-documented examples where regulation is accomplished effectively by only a few TF’s in a cell (e.g., about 10 for *lac*-repressor in *E. coli*) [24].

As studied in detail by Winter, Berg, and von Hippel [8, 9, 10], the search dynamics of TF’s involves instead a combination of sliding along the DNA at short length scales and *hopping* between different segments of DNA (either over the dissociation barrier through the cytoplasm, or by direct intersegment transfer); see Fig. 2(a). This search mode is much faster (given the high DNA concentration inside the cell), since the dynamics is essentially 3D diffusion beyond the hopping scale, and 3D diffusion is much less redundant than 1D diffusion. For example, if the TF’s were not bound to the DNA at all, a single TF of a few nm in linear dimension ℓ would locate its target in a cell volume V_{cell} of several μm^3 in the average first passage time of $T_{3D} = V_{\text{cell}}/(4\pi\ell D_3) \sim 10 \text{ sec}$, given a 3D diffusion constant on the order of $D_3 \sim 10 \mu\text{m}^2/\text{sec}$ [25]. The search time $T_{3D/1D}$ for the combined 1D/3D diffusion under *in vivo* conditions can be estimated to be comparable to T_{3D} [10]. Hence, the search time is short enough to comfortably allow even a single TF to locate its target within the physiological time scale.

In the study of the search dynamics reviewed above, binding of the TF to the genomic background was assumed to occur at a single energy value, namely, the non-specific energy ΔG_{ns} [8]. On the other hand, the ‘‘energy landscape’’ of Fig. 1(b) clearly shows that the random genomic background contains many isolated sites with binding energies far below ΔG_{ns} . These sites constitute kinetic traps which can, in principle, drastically impede the local search process, if the energy difference to their surroundings is sufficiently large⁵. Thus to fully understand the search dynamics, we need to character-

³ In Ref. [17], the nonspecific binding was not included so that $\overline{Z_b} = \overline{Z_{sp}}$, and the energy scale was shifted such that $\overline{Z_b} = N$.

⁴ In many applications, including protein folding [23], the REM was introduced to *approximate* the random background interaction. The TF-DNA interaction as defined by (5) represents one of the few systems for which the REM description is *directly applicable*.

⁵ Note that the additional sequence-specific binding energy to a ‘spurious site’ in the background equally increases the kinetic barrier for sliding to a neighboring site as well as for dissociation into the cytoplasm.




FIG. 2 (a) Schematic illustration of the search dynamics: a TF (represented by a solid ellipse) moves among genomic DNA (lines) via a combination of 1D diffusion (along the genome) and 3D diffusion (hopping between nearby segments) as illustrated by the arrows. The open circles indicate the potential kinetic traps which are sites that are preferred by the TF in a random background. (b) Dependence of the chemical potential μ on the number n of TF's in a cell for Mnt, Cro, and λ -repressor, obtained by directly solving and inverting the defining equation (13). The comparison with the dashed line $\mu = k_B T \ln n$ shows that $\mu(n)$ is sufficiently well described by the simple expression (14) over the regime $1 < n < 1000$.

ize the effect of kinetic traps in the genomic background: what is the constraint on the design of TF-DNA interaction imposed by requiring that the effect of kinetic traps be negligible?

At each binding sequence $\vec{s}_j$ with energy $E_j \equiv E[\vec{s}_j] < E_{ns}$, the TF typically spends a time $\tau_j = \tau_0 \cdot e^{\beta(E_{ns} - E[\vec{s}_j])}$, where τ_0 is the average ‘waiting time’ of the TF at a non-specific binding site. Along the search path of the TF, the *average waiting time* $\bar{\tau}$ per binding site is then simply given by

$$\bar{\tau} = \tau_0 \sum_E \Omega_{sp}(E) [e^{\beta(E_{ns} - E)} \theta(E_{ns} - E) + \theta(E - E_{ns})]. \quad (10)$$

Here we assumed as before that the genomic sequence is random so that the sequence-specific binding energy E can be treated as a random variable drawn from the distribution $\Omega_{sp}(E)$. The second term, with the help of the unit step function $\theta(x)$, is used to express the fact that there is no kinetic trap for the (majority of) sites with $E > E_{ns}$.

A comparison of (10) and Eq. (8) for the average partition function $\bar{Z}_b$ immediately yields the important relation⁶

$$\bar{\tau}/\tau_0 \approx \bar{Z}_b e^{\beta E_{ns}}/N, \quad (11)$$

since in Eq. (8), the second term dominates for $E > E_{ns}$. As expected, the kinetic trap factor $\bar{\tau}/\tau_0$ grows exponentially with E_{ns} , the threshold to nonspecific binding.

⁶ Note that this relation is actually independent of the additive form of the binding energy (2).

On the other hand, we note from $\bar{Z}_b = \bar{Z}_{sp} + N e^{-\beta E_{ns}}$ (see ‘Thermodynamics’) that the trap factor can be made to be of order one so that the dynamical analysis of Refs. [8, 9, 10] remains qualitatively valid, if $\bar{Z}_{sp} \leq N e^{-\beta E_{ns}}$. The physical meaning of this condition is that the average effect of the kinetic traps can be rendered small if the sum of the waiting times does not exceed the order of the plain diffusion time. As we will see, this can be accomplished by choosing the binding energy matrix $\mathcal{E}_i(s)$ and E_{ns} appropriately. Combining this kinetic constraint with Eq. (9), we obtain the condition

$$\bar{Z}_{sp} \leq N e^{-\beta E_{ns}} \leq 1 \quad (12)$$

for the rapid recognition of a target sequence by a single TF.

Programmability of Binding Threshold

Multiple transcription factors

There are of course typically multiple copies of the same TF in the cell, and the regulatory function is accomplished if any one of these TF's binds to the target sequence. If the cell contains n copies of a given TF, then the occupation probability for the target sequence, Eq. (3), is replaced by the Fermi distribution (or ‘Arrhenius function’) $P_t = 1/(1 + e^{\beta(E_t - \mu)})$, since each binding sequence can at most be occupied by one TF. The chemical potential $\mu(n)$ is determined implicitly from the condition⁷

$$n = \sum_E [\Omega_{sp}(E) + \delta_{E, E_{ns}}] \cdot \frac{1}{1 + e^{\beta(E - \mu)}}, \quad (13)$$

where the quantity in brackets represents the total density of states. In the simplest scenario, where steric exclusion between TF's bound to the non-target sequences is negligible, one has [11]

$$\mu \approx k_B T \ln n + F_b. \quad (14)$$

This is empirically found to be a good approximation for those TF's with known binding energy matrices, as shown in Fig. 2(b). We will adopt the form (14) for the chemical potential of a generic TF in this study; a general argument will be given later to justify this choice, even for the case where multiple target sequences are present in the same genome.

Using (14), the occupation probability can be more succinctly written as $P_t = 1/(1 + \tilde{n}_t/n)$, where

$$\tilde{n}_t = e^{\beta(E_t - F_b)} \quad (15)$$

⁷ Here, the exclusion between overlapping binding sites can be neglected since $n \ll N$. Also, we have not included the (unimportant) exclusion between the specific and the unspecific binding mode at a given site.

denotes the (soft) *threshold concentration* of the TF for occupation of the target sequence.

Programmability

The allowed values of the background free energy F_b for the binding of the target sequence obviously depend on the TF concentration n . For example, we have the condition (4) for $n = 1$, while smaller values are allowed for $n > 1$. It thus appears that the allowed F_b 's are different for the different TF's, since they would typically be present in the cell with different concentrations. On the other hand, even for a given TF species, the *desired* binding threshold may not be at a *single* concentration for different target sites, but can *vary* depending on functional demands. For example, it can be desirable to turn on different genes/operons at different TF concentrations, in order to maintain a *temporal order* in the expression of different operons as the concentration of the controlling TF gradually changes over time. This effect was observed recently for the *E. coli* flagella assembly [7] and SOS response systems (U. Alon, private communication).

As another example, consider the case where a particular TF A is involved in the regulation of two operons, X and Y. Suppose it is desired that A activates the transcription of operon X on its own at a concentration n_A , while operon Y should *only* be activated if A is present (at the same concentration n_A) *together* with another TF B that can bind cooperatively with A. It is then desirable to have a strong binding site for A in the regulatory region of operon X such that its threshold $\tilde{n}_{A,X} < n_A$, and a weak binding site in the regulatory region of operon Y, with a threshold $\tilde{n}_{A,Y} > n_A$. The latter insures that the operon Y will not be accidentally activated by fluctuations in n_A alone, and only when the TF B is present would the attractive interaction between A and B induce the two to bind to their targets.

The above examples show that it is functionally desirable to have the ability to *set* the binding threshold $\tilde{n}_t$ of a given TF to each of its target sequence $\vec{s}_t$ *individually*. As is clear from the defining expression (15), this can only be done through the choice of the target sequence $\vec{s}_t$ which affects E_t , since the other variable, F_b , is fixed for a given TF. We refer to the ability to control the binding threshold $\tilde{n}_t$ through the choice of the target sequence $\vec{s}_t$ *alone* as “programmability” of the binding threshold. Assuming that programmability is a desirable feature of TF-DNA interaction (since sequence changes can be easily accomplished by point mutation if the functional need arises), we seek to determine the specifics of the TF-DNA interaction, e.g., the binding matrix $\mathcal{E}_i(s)$, the length of the binding sequence L , and the threshold energy E_{ns} , which allow the targets to be *maximally programmable*.

Two-state model and parameter selection

Specifically, let us require programmability of the binding threshold over the entire range $\tilde{n} = 1 \dots 10^3$, since typical cellular TF concentrations range from a few to a few hundred per cell. The lower bound $\tilde{n} \approx 1$ immediately imposes the condition (4) on F_b , or, taking also the kinetic constraint into account, the condition (12). Furthermore, in order to tune $\tilde{n}$ throughout the desired range with a reasonable resolution, it is necessary to have the ability to change E_t from 0 to $k_B T \ln 10^3 \approx 7k_B T$, in small increments. This requires the non-zero entries of the binding energy matrix $\mathcal{E}_i(s)$ to take on small values. Which choices for the TF-DNA interaction parameters ($\mathcal{E}_i(s)$, L , E_{ns}) can simultaneously satisfy the latter requirement and condition (12)?

The combined effect of these physical constraints and functional demands is best understood by simplifying the energy matrix $\mathcal{E}$ such that we retain the essential and generic aspect of sequence-specific binding, while eliminating all TF-specific details. Towards this end, we adopt the two-state model originally introduced by von Hippel and Berg [11], characterizing all of the non-zero entries of the *significant positions*⁸ in the energy matrix by a *single* value, i.e.,

$$\beta\mathcal{E}_i(s) = \begin{cases} 0 & \text{if } s = s_i^* \\ \varepsilon > 0 & \text{if } s \neq s_i^* \end{cases}, \quad (16)$$

where ε is a dimensionless “discrimination energy” (in units of $k_B T$). It describes the energetic *preference* of the TF for the optimal binding sequence $\vec{s}^*$, and is a crucial parameter controlling the specificity of the TF. Within the two-state model, the binding energy to the target $\vec{s}_t$ is simply ε times the total number of mismatches between the target and the best binder $\vec{s}^*$, i.e., $E[\vec{s}_t] = \varepsilon \cdot |\vec{s}_t - \vec{s}^*|$, where $|\dots|$ denotes the Hamming distance between two sequences. Clearly, programmability is best satisfied with a small ε which enhances the resolution of the programmable binding threshold.

The two-state model (16) also allows an explicit evaluation of the condition (12) via the formulae (6) for $\overline{Z}_{sp}$. Assuming for simplicity equal single-nucleotide frequencies in the background (i.e., $p(s) = 1/4$), the quantity in the bracket of (6) is easily evaluated. We have $\overline{Z}_{sp}(\varepsilon, L) = N \cdot \zeta^L(\varepsilon)$ where $\zeta \equiv \sum_s e^{-\beta\mathcal{E}(s)} p(s) = (1 + 3e^{-\varepsilon})/4$. Note that ζ^{-1} is in the range between 1 and 4, and can be regarded as the effective size of the nucleotide “alphabet” as “seen” by the TF in the specific binding mode. The maximum value $\zeta^{-1} = 4$ is attained if the energy matrix has infinite discrimination, $\varepsilon \rightarrow \infty$, while no discrimination can be achieved at $\varepsilon = 0$ where $\zeta^{-1} = 1$. In

⁸ Note that the energy matrices for most TF's contain a number of (fixed) positions which have no strong preference for any of the nucleotides. We will not consider these positions in the ensuing discussion of the two-state model, and will use L to refer to the total number of significant positions.




FIG. 3 (a) Plot of the region where $\overline{Z_{sp}}(\varepsilon, L) \leq 1$. The boundary $L^*(\varepsilon)$ for $N = 10^7$ is indicated by the solid line; see text. The dashed line $\ln(N)/(\ln \zeta^{-1}(\varepsilon) - \varepsilon/(1 + e^\varepsilon/3))$ indicates the onset of the glass transition in the random energy model where the annealed approximation breaks down. As argued in the text, the desired parameter regime is close to $\overline{Z_{sp}} = 1$, so that the annealed approximation is justified. (b) The binding threshold $\tilde{n}$ as a function of the total number of mismatches r of the target sequence $\vec{s}_t$ from the best binder $\vec{s}^*$ at different parameter combinations (ε, L) .

Fig. 3(a), we indicate the allowed region $\overline{Z_{sp}}(\varepsilon, L) \leq 1$ in the parameter space of (ε, L) , with the boundary $L^*(\varepsilon) = \ln N / \ln \zeta^{-1}(\varepsilon)$ defined by $\overline{Z_{sp}}(L^*, \varepsilon) = 1$. From the figure, it is clear that the desire for small ε pushes the system to the boundary at $\overline{Z_{sp}} = 1$. Along the boundary, the smallest ε is given by the largest allowable binding length L . For typical bacterial TF's whose binding sequences are no longer than about 15 bp (usually dimers), we find $\varepsilon \approx 2$.

While the result on ε is somewhat specific to the two-state model, the need for $\overline{Z_{sp}} \rightarrow 1$ imposed by the programmability consideration forces the threshold energy to take on the value

$$E_{ns} = k_B T \ln N \approx 16 k_B T \quad (17)$$

(for $N \sim 10^7$) according to the condition (12) independent of the specifics of the binding energy matrix $\mathcal{E}$. It also follows that

$$F_b \approx 0 \quad (18)$$

so that the binding threshold is simply given by

$$\tilde{n}[\vec{s}_t] \approx e^{\beta E[\vec{s}_t]}. \quad (19)$$

The dependences of the $\tilde{n}$ on the number of mismatches for the two-state model are shown in Fig. 3(b). We see that at the optimal parameter choice of $(\varepsilon = 2, L = 15)$, each mismatch increases the binding threshold $\tilde{n}$ by nearly 10-fold. In principle, further fine-tuning can be accomplished by utilizing small variations in the mismatch energies.

Discussion

The key results of this study, that maximal programmability of the binding threshold $\tilde{n}$ requires the TF-DNA interaction to satisfy the conditions (17) and (18),




FIG. 4 Graphical construction of the background free energy F_b and other quantities used in the text.

can be conveniently summarized graphically using the density of states $\Omega_{sp}(E)$. In Fig. 4, the density of states is plotted with the normalization that $\max_E \Omega_{sp}(E) = N$, as indicated by the horizontal dotted line. The background free energy F_b can be obtained using the Legendre construction: One draws the line $e^{\beta(E - F_b)}$ (the dashed line in the semi-log plot of Fig. 4) such that it just touches $\Omega_{sp}(E)$. F_b can then be read off as the intercept of the dashed line on the E -axis, which should be in the vicinity of the origin according to (18). Similarly, E_{ns} [as given by (17)] can be read off as the E -coordinate where the dashed line intersects the horizontal dotted line.

The point where the dashed line tangents $\Omega_{sp}(E)$ is also physically meaningful: The E -coordinate of the tangent point gives the ensemble-averaged binding energy $E_0 \equiv \sum_E E \Omega_{sp}(E) e^{-\beta E} / Z_{sp}$. The vertical coordinate N_0 of the tangent point is given by the relation $F_b = E_0 - k_B T \ln N_0$, which expresses the fact that the dominant contribution to the background free energy stems from the N_0 sequences of energy $\approx E_0$ in the collection of N random sequences: The Boltzmann weight of those sequences with $E > E_0$ is too small to contribute to the partition sum, while for $E < E_0$, there are too few sequences.

The value of N_0 is an important characteristics of the system. $S = \ln N_0$ is known as the “entropy” of this system, and $H = \ln(N/N_0)$ is known as the “relative entropy”; the latter has been used to characterize the specificity of the TF-DNA interaction [17]. As mentioned before, the annealed approximation is only valid if many terms contribute to the partition sum, i.e., if $N_0 \gg 1$ or $S > 0$. For the two-state model (16), the values of ε and L corresponding to the line $N_0 = 1$ is far from the line $L^*(\varepsilon)$ selected by the maximal programmability criterion; this justifies the use of the annealed approximation. At the optimal parameter of $\varepsilon = 2$ and $L = 15$, we have $N_0 \approx 10^3 \gg 1$. The corresponding relative entropy is $H \approx 7$ (about 10 bits).

The large value of N_0 also provides us with an intuitive understanding of the simple dependence (14) of the chemical potential μ on the cellular TF concentration n ; see Fig. 2(b). As mentioned already, the expression (14) is obtained if multiple occupancy of the background

	theory	<i>Mnt</i>	<i>Cro</i>	<i>cI</i>	<i>LacR</i>
F_b [$k_B T$]	0	-1.2	-1.6	-0.8	–
H [bits]	$\approx$ 10	8.9	13.5	12.7	–
E_{ns} [$k_B T$]	16	$17^\dagger$	–	–	≈ 16

TABLE I Comparison of the expected values of the background free energy F_b , relative entropy H and the threshold to nonspecific binding E_{ns} to the known values of these parameters for *Mnt*, *Cro*, the λ -repressor *cI*, and the *lac*-repressor *LacR*. The units of these values are given by the bracket in the first column. A dash indicates that the value is not available. [†]see Ref. [28]

sequences is negligible at the TF concentration n . Since there is a large number (i.e., N_0) binding sequences which contribute significantly to the net effect of background binding, multiple occupancy of these sequences is indeed not likely if $n < N_0$. Thus for $N_0 \sim O(10^3)$, the expression (14) can be taken as a good approximation of the chemical potential over the typical range of cellular TF concentration $n = 1 \dots 10^3$, as shown in Fig. 2(b) for the 3 known TF's. We expect this result to hold even if there are multiple target sequences, say m_t , whose binding energy E_t is much lower than E_0 , as long as $E_t > k_B T \ln m_t$ such that F_b is not affected by the addition of these target sequences to the density of states. Having $\mu(n)$ independent of the number of targets is a desirable functional robustness property from a system perspective, since one wouldn't want to perturb the recognition of the TF's and the existing targets by the addition of a few new targets. It will be interesting to see to what extent this feature is preserved by studying the energetics of TF's with a large number of target sites, e.g., the catabolic repressor protein CRP in *E. coli* [5].

Finally, we compare the values of the optimal interaction parameters according to our theory to those of the well-studied TF's. From the values listed in Table I, we see that all of the available data is in the neighborhood of the expectation based on the maximal programmability criterion. We do not suggest here that programmability was necessarily the selective driving force that constrained the TF-DNA interaction to its observed form (there could be other reasons, e.g. biochemical restrictions, for the interaction to be of this form). However, the rough correspondence between theory and observation does indicate that it is *possible* (and perhaps even very likely) that TF's generally have the required energetics for their binding threshold to be programmable over a wide range.

One obvious short-coming of the above comparison is that the three TF's for which the interaction parameters are known are all from bacteriophages and may not represent typical prokaryotic TF's. It will therefore be very important to experimentally determine the interaction parameters for a variety of different TF's. The results of a sufficient number of such studies will inform us whether programmability is a generic feature of TF-DNA interaction. Knowledge of this kind can be very helpful

in developing appropriate coarse-grained models of gene regulation at the system level. In particular, quantitative relations of the type suggested by Eq. (19) will be necessary for an eventual quantitative description of gene regulatory networks. Also, this knowledge would have important implications for the evolution of gene regulation [26, 27].

Acknowledgments. — It is a pleasure to acknowledge useful discussions with G. Stormo, P. von Hippel, and K. Sneppen on many aspects of TF-DNA interaction. We are also grateful to the hospitality of the Institute for Theoretical Physics in Santa Barbara where some of the work was carried out. This research is supported in part by the National Science Foundation through Grant Nos. DMR-9971456. UG was supported in part by a fellowship from the 'Deutscher Akademischer Austauschdienst', and TH by a Burroughs-Wellcome functional genomics award.

References

- [1] Davidson, E.H. *et al.* (2002) *Science* **295**, 1669–1677.
- [2] Berman, B.P. *et al.* (2002) *Proc. Natl. Acad. Sci. USA* **99**, 757–762.
- [3] Bussemaker, H.J., Li, H. & Siggia, E.D. (2001) *Nature Genet.* **27**, 167–171.
- [4] Yuh, C., Bolouri, H. & Davidson, E.H. (2001) *Development* **128**, 617–629.
- [5] Neidhardt, F.C., ed. (1996) *Escherichia coli and Salmonella: cellular and molecular biology* (ASM Press, Washington D.C.)
- [6] Ptashne, M. & Gann, A. (2002) *Genes & Signals* (Cold Spring Harbor Laboratory Press, Cold Spring Harbor, N.Y.)
- [7] Kalir, S., McClure, J., Pabbaraju, K., Southward, C., Ronen, M., Leibler, S., Surette, M.G. & Alon, U. (2001) *Science* **292**, 2080–2083.
- [8] Berg, O.G., Winter, R.B. & von Hippel, P.H. (1981) *Biochemistry* **20**, 6929–6948.
- [9] Winter, R.B. & von Hippel, P.H. (1981) *Biochemistry* **20**, 6948–6960.
- [10] Winter, R.B., Berg, O.G. & von Hippel, P.H. (1981) *Biochemistry* **20**, 6961–6977.
- [11] von Hippel, P.H. & Berg, O.G. (1986) *Proc. Natl. Acad. Sci. USA* **83**, 1608–1612.
- [12] Berg, O.G. & von Hippel, P.H. (1987) *J. Mol. Biol.* **193**, 723–750.
- [13] deHaseth, P.L., Gross, C.A., Burgess, R.R. & Record, M.T. Jr. (1977) *Biochemistry* **16**, 4777–4783.
- [14] Record, M.T., Jr., deHaseth, P.L. & Lohman, T.M. (1977) *Biochemistry* **16**, 4791–4796.
- [15] Kao-Huang, Y., Revzin, A., Butler, A.P., O'Conner, P., Noble, D.W. & von Hippel, P.H. (1977) *Proc. Natl. Acad. Sci. USA* **74**, 4228–4232.
- [16] Fields, D.S., He, Y., Al-Uzri, A.Y. & Stormo, G.D. (1997) *J. Mol. Biol.* **271**, 178–194.
- [17] Stormo, G.D. & Fields, D.S. (1998) *Trends in Biochemical Sciences* **23**, 109–113.
- [18] Sarai, A. & Takeda, Y. (1989) *Proc. Natl. Acad. Sci. USA* **86**, 6513–6517.
- [19] Takeda, Y., Sarai, A. & Rivera, V.M. (1989) *Proc. Natl. Acad. Sci. USA* **86**, 439–443.

- [20] Frank, D.E., Saecker, R.M., Bond, J.P., Capp, M.W., Tsodikov, O.V., Melcher, S.E., Levandoski, M.M. & Record, M.T., Jr. (1997) *J. Mol. Biol.* **267**, 1186–1206.
- [21] Pabo, C.O. & Sauer, R.T. (1992) *Annu. Rev. Biochem.* **61**, 1053–1095.
- [22] Derrida, B. (1981) *Phys. Rev. B* **24**, 2613–2626.
- [23] Bryngelson, J.D. & Wolynes, P.G. (1987) *Proc. Natl. Acad. Sci. USA* **84**, 7524–7528.
- [24] Droge, P. & Muller-Hill, B. (2001) *Bioessays* **23**, 179–183.
- [25] Elowitz, M.B., Surette, G.S., Wolf, P.-E., Stock, J.B. & Leibler, S. (1999) *J. Bacteriol.* **181**, 197–203.
- [26] Sengupta, A.M., Djordjevic, M. & Shraiman, B.I. (2002) *Proc. Natl. Acad. Sci. USA* **99**, 2072–2077.
- [27] Gerland, U. & Hwa, T. (2002) *J. Mol. Evol.* **55**, 386–400.
- [28] Raumann, B.E., Knight, K.L. & Sauer, R.T. (1995) *Nat. Struct. Biol.* **2**, 1115–1122.